



Cell Adhesion and Migration

Antibodies for blocking and detecting proteins involved in cell migration and adhesion

bioxcell.com



Cell Adhesion and Migration Antibodies



InVivoMab vs. InVivoPlus		
	InVivoMab > 95%	InVivoPlus > 95%
purity level	> 95%	> 95%
protein aggregates validated at ≤ 5%	✓	✓
azide and carrier protein free	✓	✓
endotoxin concentration	< 2EU/mg	< 1EU/mg
validated by immunoblot, FC, or ELISA	✓	✓
tested for murine pathogens	✓	✓
available in bulk quantities	✓	✓

Antigen	Reactivity	Application	Clone	InVivoMab Catalog	InVivoPlus Catalog
CD11a (LFA-1α)	human	Functional assays, FC	R7-1	BE0192	
CD18	mouse	<i>in vivo</i> LFA-1 neutralization	M18/2	BE0009	
CD22	mouse	<i>in vivo</i> B cell depletion in combination with anti-CD19 (clone 1D3) and anti-rat κ Light Chain (clone MAR 18.5), FC, IP	Cy34.1	BE0011	
CD24	mouse	<i>in vivo</i> administration, IHC-F, IHC-P, IF, FC	M1/69	BE0360	
CD29	mouse	IF, FC	KMI6	BE0232	
CD44	mouse/human	<i>in vivo</i> CD44 neutralization	IM7	BE0039	
CD44	human	<i>in vivo</i> CD44 blockade in xenografts, <i>in vitro</i> CD44 blockade, WB, IF	Hermes-1	BE0262	
CD47	human	<i>in vitro</i> CD47 neutralization, <i>in vivo</i> CD47 neutralization in human tumor xenograft models or humanized mice, FC	B6H12	BE0019-1	
CD47	human/mouse/rat	<i>in vivo</i> and <i>in vitro</i> CD47 blockade, IF	MIAP410	BE0283	BP0283
CD47 (IAP)	mouse	<i>in vivo</i> CD47 blockade, <i>in vitro</i> CD47 blockade, IF	MIAP301	BE0270	
CD54 (ICAM-1)	mouse	<i>in vivo</i> ICAM-1 neutralization	YN1/1.7.4	BE0020-1	
CD54 (ICAM-1)	human	<i>in vitro</i> T cell stimulation/activation, IF	R6-5-D6	BE0020-2	
CD96	mouse	<i>in vivo</i> and <i>in vitro</i> CD96 blocking, FC	3.3	BE0337	
CD103	mouse	<i>in vivo</i> CD103 neutralization, IF, FC	M290	BE0026	
CD106 (VCAM-1)	mouse	<i>in vivo</i> VCAM-1 neutralization, IF	M/K-2.7	BE0027	
CD172a (SIRPα)	mouse	<i>in vivo</i> and <i>in vitro</i> SIRPα blocking, WB, IP, FC	P84	BE0322	
CD209b (SIGN-R1)	mouse	<i>in vivo</i> SIGN-R1 blockade, IHC-F, WB, FC	22D1	BE0220	
CD317 (BST2, PDCA-1)	mouse	<i>in vivo</i> pDC depletion, IF, FC	927	BE0311	
CEACAM 1/3/5/6/8	human	WB, ELISA, IF, IHC-P, FC	6G5j	BE0362	
CXCL9 (MIG)	mouse	IF	MIG-2F5.5	BE0309	
E-Cadherin (CD324)	mouse	<i>in vivo</i> E-Cadherin neutralization, <i>in vitro</i> E-Cadherin neutralization, IF, IP, WB	DECMA-1	BE0352	
E-Selectin	human	FC	CL2	BE0136	
E-selectin (CD62E)	mouse	<i>in vivo</i> and <i>in vitro</i> E-selectin blockade, IHC-F	9A9	BE0294	
Galectin-9	mouse	<i>in vivo</i> and <i>in vitro</i> Galectin-9 blockade	RG9-1	BE0218	
IL-1α	mouse	<i>in vivo</i> IL-1α neutralization	ALF-161	BE0243	
IL-1β	mouse/rat	<i>in vivo</i> and <i>in vitro</i> IL-1β neutralization, ELISA	B122	BE0246	
Integrin β7	mouse/human	<i>in vivo</i> blocking of integrin β7, IP, WB, FC	FIB504	BE0062	
LFA-1α (CD11a)	human	<i>in vitro</i> LFA-1 neutralization	TS-1/22.1.13	BE0005	
LFA-1α (CD11a)	mouse	<i>in vivo</i> LFA-1 neutralization	FD441.8	BE0005-1	
LFA-1α (CD11a)	mouse	<i>in vivo</i> LFA-1 neutralization, FC	M17/4	BE0006	
LPAM-1 (Integrin α4β7)	mouse	<i>in vivo</i> Integrin α4β7 neutralization, FC	DATK32	BE0034	
LRP1 (CD91)	mouse/human/rat	WB, IF, IP	11H4	BE0333	
L-Selectin (CD62L)	mouse	<i>in vivo</i> CD62L neutralization	Mel-14	BE0021	
MAdCAM-1	mouse	<i>in vivo</i> MAdCAM-1 neutralization, IF	MECA-367	BE0035	
MUC1 (CD227)	human	<i>in vivo</i> administration in mouse xenograft models, IHC-P, IF, <i>in vitro</i> cell cytotoxicity assay, WB	C595 (NCRC48)	BE0336	
PSGL-1 (CD162)	mouse	<i>in vivo</i> PSGL-1 blockade, IHC-F	4RA10	BE0186	
Siglec-H	mouse	<i>in vivo</i> administration, FC	440c	BE0202	
Thy1.1 (CD90.1)	mouse	<i>in vivo</i> T cell depletion	19E12	BE0214	
Thy1.2 (CD90.2)	mouse	<i>in vivo</i> T cell depletion, <i>in vivo</i> ILC depletion, WB	30H12	BE0066	BP0066
VEGFR-2	mouse	<i>in vivo</i> blocking of VEGF/VEGFR-2 signaling, <i>in vitro</i> blocking of VEGFR signaling	DC101	BE0060	BP0060
VLA-4 (CD49d)	mouse/human	<i>in vivo</i> and <i>in vitro</i> VLA-4 neutralization, FC	PS/2	BE0071	

IF Immunofluorescence | IHC-F Immunohistochemistry (frozen) | IHC-P Immunohistochemistry (paraffin) | WB Western blot | IP Immunoprecipitation | FC Flow cytometry (requires fluochrome conjugation)

For over 20 years, scientists have trusted Bio X Cell as their go-to source for *in vivo* functional grade antibodies. This is reflected in over 15,000 peer-reviewed publications citing Bio X Cell products. We understand this responsibility is of paramount importance and remain committed to producing antibodies of unparalleled quality and consistency, enabling our partners around the globe to accelerate research and discoveries.

Bio X Cell
 bioxcell.com | 1.866.787.3444 | customerservice@bioxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell. © Bio X Cell 2021

